

Post Operative Care (Wound Care)

Day 1:

- 1. Leave the bandage on for 24 hours.
- 2. Limit activity for the first 48 hours.
- 3. If you experience pain, you may alternate ibuprofen and acetaminophen every 3 hours unless contraindicated. Do not take aspirin as this may increase bleeding.

Day 2 and following:

- 1. Wash your hands and remove the bandage.
- 2. Gently clean the suture site with soap and water.
- 3. Blot the site dry with a clean soft cloth.
- 4. Apply Vaseline or Aquaphor using a q-tip. It is important to keep the site moist with ointment to avoid scab formation. This aids in the healing process and reduces scarring.
- 5. Apply a clean bandage.
- 6. Repeat process once daily.
- 7. Showering is acceptable, but avoid prolonged submersion of the site in water. If shaving, be careful not to shave over suture area.

noval or	' wound	cneck
ľ	noval or	noval or wound

What to do if:

- Bleeding:
 - Apply firm pressure to site for 10 minutes. If the bleeding does not stop, please call the clinic.
 - To prevent bleeding, avoid activity and straining. Keep the site elevated.

What to watch for:

- Some redness, swelling and drainage at the surgical site are normal.
 These are expected. Signs of infection include increased pain or yellow/green drainage from the site.
- If you notice any of these signs, call the clinic at (970)247-1970.

523 S. Camino del Rio Durango, CO 81303 (970) 247-1970 www.durangodermatology.com